arXiv:1606.03872v2 [math.CO] 17 Jun 2016

The (k, ℓ)-proper index of graphs*

Hong Chang[1], Xueliang Li[1], Colton Magnant[2], Zhongmei Qin[1]

[1]Center for Combinatorics and LPMC

Nankai University, Tianjin 300071, P.R. China

Email: changh@mail.nankai.edu.cn, lxl@nankai.edu.cn, qinzhongmei90@163.com

[2]Department of Mathematical Sciences

Georgia Southern University, Statesboro, GA 30460-8093, USA

Email: cmagnant@georgiasouthern.edu

Abstract

A tree T in an edge-colored graph is called a *proper tree* if no two adjacent edges of T receive the same color. Let G be a connected graph of order n and k be an integer with $2 \leq k \leq n$. For $S \subseteq V(G)$ and $|S| \geq 2$, an S-tree is a tree containing the vertices of S in G. Suppose $\{T_1, T_2, \ldots, T_\ell\}$ is a set of S-trees, they are called *internally disjoint* if $E(T_i) \cap E(T_j) = \emptyset$ and $V(T_i) \cap V(T_j) = S$ for $1 \leq i \neq j \leq \ell$. For a set S of k vertices of G, the maximum number of internally disjoint S-trees in G is denoted by $\kappa(S)$. The κ-connectivity $\kappa_k(G)$ of G is defined by $\kappa_k(G) = \min\{\kappa(S) \mid S \text{ is a } k\text{-subset of } V(G)\}$. For a connected graph G of order n and for two integers k and ℓ with $2 \leq k \leq n$ and $1 \leq \ell \leq \kappa_k(G)$, the (k, ℓ)-*proper index* $px_{k,\ell}(G)$ of G is the minimum number of colors that are needed in an edge-coloring of G such that for every k-subset S of $V(G)$, there exist ℓ internally disjoint proper S-trees connecting them. In this paper, we show that for every pair of positive integers k and ℓ with $k \geq 3$, there exists a positive integer $N_1 = N_1(k, \ell)$ such that $px_{k,\ell}(K_n) = 2$ for every integer $n \geq N_1$, and also there exists a positive integer $N_2 = N_2(k, \ell)$ such that $px_{k,\ell}(K_{m,n}) = 2$ for every integer $n \geq N_2$ and $m = O(n^r)(r \geq 1)$. In addition, we show that for every $p \geq c\sqrt[k]{\frac{\log_a n}{n}}$ $(c \geq 5)$, $px_{k,\ell}(G_{n,p}) \leq 2$ holds almost surely, where $G_{n,p}$ is the Erdös-Rényi random graph model.

Keywords: proper tree; proper index; random graphs; threshold function.

*Supported by NSFC No.11371205, 11531011, "973" program No.2013CB834204 and PCSIRT.

AMS subject classification 2010: 05C15, 05C40, 05C80, 05D40.

1 Introduction

All graphs in this paper are undirected, finite and simple. We follow [4] for graph theoretical notation and terminology not described here. Let G be a nontrivial connected graph with an associated *edge-coloring* $c : E(G) \to \{1, 2, \ldots, r\}$, $r \in \mathbb{N}$, where adjacent edges may have the same color. If adjacent edges of G are assigned different colors by c, then c is a *proper coloring*. The minimum number of colors needed in a proper coloring of G is referred to as the *chromatic index* of G and denoted by $\chi'(G)$. A path of G is said to be a *rainbow path* if no two edges on the path receive the same color. The graph G is called *rainbow connected* if for every pair of distinct vertices there is a rainbow path of G connecting them. An edge-coloring of a connected graph is a *rainbow connecting coloring* if it makes the graph rainbow connected. This concept of rainbow connection of graphs was introduced by Chartrand et al. [12] in 2008. The *rainbow connection number* $rc(G)$ of a connected graph G is the smallest number of colors that are needed in order to make G rainbow connected. The readers who are interested in this topic can see [21, 22] for a survey.

In [13], Chartrand et al. generalized the concept of rainbow connection to rainbow index. At first, we recall the concept of generalized connectivity. Let G be a connected graph of order n and k be an integer with $2 \leq k \leq n$. For $S \subseteq V(G)$ and $|S| \geq 2$, an S-tree is a tree containing the vertices of S in G. Let $\{T_1, T_2, \ldots, T_\ell\}$ be a set of S-trees, they are called *internally disjoint* if $E(T_i) \cap E(T_j) = \emptyset$ and $V(T_i) \cap V(T_j) = S$ for every pair of distinct integers i, j with $1 \leq i, j \leq \ell$. For a set S of k vertices of G, the maximum number of internally disjoint S-trees in G is denoted by $\kappa(S)$. The κ-connectivity $\kappa_k(G)$ of G is defined by $\kappa_k(G) = \min\{\kappa(S) \mid S \text{ is a } k\text{-subset of } V(G)\}$. We refer to a book [20] for more details about the generalized connectivity.

A tree T in an edge-colored graph is a *rainbow tree* if no two edges of T have the same color. Let G be a connected graph of order n and let k, ℓ be two positive integers with $2 \leq k \leq n$ and $1 \leq \ell \leq \kappa_k(G)$. The (k, ℓ)-*rainbow index* of a connected graph G, denoted by $rx_{k,\ell}(G)$, is the minimum number of colors that are needed in an edge-coloring of G such that for every k-subset S of $V(G)$, there exist ℓ internally disjoint rainbow S-trees connecting them. Recently, a lot of relevant results have been published in [6, 7, 8]. In particular, for $\ell = 1$, we write $rx_k(G)$ for $rx_{k,1}(G)$ and call it the k-*rainbow index* of G (see [9, 10, 15]).

Motivated by rainbow coloring and proper coloring in graphs, Andrews et al. [2] and Borozan et al. [5] introduced the concept of proper-path coloring. Let G be a nontrivial connected graph with an edge-coloring. A path in G is called a *proper path* if no two adjacent edges of the path are colored with the same color. An edge-colored graph G is *proper connected* if any two vertices of G are connected by a proper path. For a connected graph G, the *proper connection number* of G, denoted by $pc(G)$, is defined as the smallest number of colors that are needed in order to make G proper connected. The *k-proper connection number* of a connected graph G, denoted by $pc_k(G)$, is the minimum number of colors that are needed in an edge-coloring of G such that for every two distinct vertices of G are connected by k internally pairwise vertex-disjoint proper paths. For more details, we refer to [17, 23] and a dynamic survey [19].

Recently, Chen et al. [14] introduced the concept of k-proper index of a connected graph G. A tree T in an edge-colored graph is a *proper tree* if no two adjacent edges of T receive the same color. Let G be a connected graph of order n and k be a fixed integer with $2 \leq k \leq n$. An edge-coloring of G is called a *k-proper coloring* if for every k-subset S of $V(G)$, there exists a proper S-tree in G. For a connected graph G, the *k-proper index* of G, denoted by $px_k(G)$, is defined as the minimum number of colors that are needed in a k-proper coloring of G. In [11], we gave some upper bounds for the 3-proper index of graphs.

A natural idea is to introduce the concept of the (k, ℓ)-proper index. Let G be a nontrivial connected graph of order n and size m. Given two integers k, ℓ with $2 \leq k \leq n$ and $1 \leq \ell \leq \kappa_k(G)$, the *$(k, \ell)$-proper index* of a connected graph G, denoted by $px_{k,\ell}(G)$, is the minimum number of colors that are needed in an edge-coloring of G such that for every k-subset S of $V(G)$, there exist ℓ internally disjoint proper S-trees connecting them. From the definition, it follows that

$$1 \leq px_{k,\ell}(G) \leq \min\{rx_{k,\ell(G)}, \chi'(G)\} \leq m.$$

Clearly, $px_{2,\ell}(G) = pc_\ell(G)$, $px_{k,1}(G) = px_k(G)$.

Let us give an overview of the rest of this paper. In Section 2, we study the (k, ℓ)-proper index of complete graphs using two distinct methods. We show that there exists a positive integer $N_1 = N_1(k, \ell)$, such that $px_{k,\ell}(K_n) = 2$ for every integer $n \geq N_1$. In Section 3, we turn to investigate the (k, ℓ)-proper index of complete bipartite graphs by probabilistic method [1]. Similarly, we prove that there exists a positive integer $N_2 = N_2(k, \ell)$, such that $px_{k,\ell}(K_{n,n}) = 2$ for every integer $n \geq N_2$. Furthermore, we can extend the result about $K_{n,n}$ to more general complete bipartite graph $K_{m,n}$, where $m = O(n^r)$, $r \in \mathbb{R}$ and $r \geq 1$. In section 4, we show that for

every $p \geq c\sqrt[k]{\frac{\log_a n}{n}}$ $(c \geq 5)$, $px_{k,\ell}(G_{n,p}) \leq 2$ holds almost surely, where $G_{n,p}$ is the Erdös-Rényi random graph model [16].

2 Complete graphs

In this section, we will investigate the (k,ℓ)-proper index of complete graphs. Firstly, we state a useful result about the k-connectivity of K_n and present some preliminary results.

Theorem 2.1 *[13] For every two integers n and k with $2 \leq k \leq n$, $\kappa_k(K_n) = n - \lceil \frac{k}{2} \rceil$.*

Theorem 2.2 *[14] Let $G = K_n$ and k be an integer with $3 \leq k \leq n$. Then $px_{k,1}(G) = px_k(G) = 2$.*

Theorem 2.3 *[5] Let $G = K_n$, $n \geq 4$ and $\ell \geq 2$. If $n \geq 2\ell$, then $px_{2,\ell}(G) = pc_\ell(G) = 2$.*

Based on the previous results, we prove the following.

Theorem 2.4 *For every integer $n \geq 4$, $px_{3,2}(K_n) = 2$.*

Proof. **Case 1.** $n = 2p$ for $p \geq 2$.

Take a Hamiltonian cycle $C = v_1, v_2, \ldots, v_{2p}$ of K_n and denote the $v_s v_t$-path in clockwise direction contained in C by $C[v_s, v_t]$. Next, we will provide an edge-coloring of K_n with 2 colors such that for any three vertices of K_n, there are two internally disjoint proper trees connecting them. We alternately color the edges of C with colors 1 and 2 starting with color 2, and color the rest of the edges with color 1. Let S be any 3-subset of $V(K_n)$, without loss of generality, we assume that $S = \{v_i, v_j, v_h\}$ with $1 \leq i < j < h \leq 2p$. It is easy to see that the Hamiltonian cycle C is partitioned into three segments $C[v_i, v_j]$, $C[v_j, v_h]$ and $C[v_h, v_i]$. If $2p+i-h = 1$, then we get that $i = 1$ and $h = 2p$. Note that the edges v_1v_2 and $v_{2p-1}v_{2p}$ are colored with color 2 and the edges $v_{2p}v_1$ and $v_{2p-1}v_1$ are colored with color 1. Thus, there are two internally disjoint proper S-trees $P_1 = v_{2p}v_1C[v_1, v_j]$ and $P_2 = v_jv_{2p}v_{2p-1}v_1$ (if $v_j = v_{2p-1}$, then $P_2 = v_{2p}v_{2p-1}v_1$). Now we suppose that $2p+i-h \geq 2$. If the edge incident to v_i with color 2 is v_iv_{i+1}, then the two internally disjoint proper S-trees are $v_hv_iC[v_i, v_j]$ and $C[v_j, v_h]C[v_h, v_i]$. Otherwise, the edge incident to v_i with color 2 is v_iv_{i-1}. Thus, the two internally disjoint proper S-trees are $v_hv_iv_{i-1}v_j$ and $C[v_i, v_j]C[v_j, v_h]$.

Case 2. $n = 2p + 1$ for $p \geq 2$.

Let H be a complete subgraph of K_n with vertex set $\{v_1, v_2, \ldots, v_{2p}\}$. Color the edges of H as above, and color the edge $v_{2p+1}v_i$ $(1 \leq i \leq 2p)$ with color 1 for i odd and with color 2 for i even. It is easy to see that for any three vertices of $V(H)$, there are two internally disjoint proper trees connecting them. Now we assume that $S = \{v_i, v_j, v_{2p+1}\}$ $(1 \leq i < j \leq 2p)$. Since there are two internally disjoint proper paths connecting v_i and v_j in the Hamiltonian cycle $C = v_1, v_2, \ldots, v_{2p}$, it follows that there are two internally disjoint proper S-trees in K_n. □

Theorem 2.5 *For every integer $n \geq 4$, $px_{n-1,2}(K_n) = 2$.*

Proof. It is well known that if n is even, then K_n can be factored into $\frac{n}{2}$ Hamiltonian paths $\{P_1, P_2, \ldots, P_{\frac{n}{2}}\}$; if n is odd, then K_n can be factored into $\frac{n-1}{2}$ Hamiltonian cycles $\{C_1, C_2, \ldots, C_{\frac{n-1}{2}}\}$. Let S be any $(n-1)$-subset of $V(K_n)$. Without loss of generality, assume that $\{v\} = V(K_n) \setminus S$. If n is even, then for each Hamiltonian path P_i, we alternately color the edges of P_i with color 1 and 2. Notice that each vertex will be an end-vertex of a Hamiltonian path. Let P_i be the Hamiltonian path which contains v as one of its end-vertices. Thus, there are two internally disjoint proper S-trees $P_i - v$ and P_j $(i \neq j)$. If n is odd, then for each Hamiltonian cycle C_i, we alternately color the edges of C_i with color 1 and 2 such that the edges incident to v_1 in each cycle are colored the same. If $v = v_1$, then there are two internally disjoint proper S-trees $C_1 - v_1$ and $C_2 - v_1$. Now we suppose $v \neq v_1$. Let C_i be the Hamiltonian cycle in which v is adjacent to v_1, and let v' be one of the neighbours of v_1 in another Hamiltonian cycle $C_j (i \neq j)$. Thus, there are two internally disjoint proper S-trees $C_i - v$ and $C_j - v_1v'$. Hence, $px_{n-1,2}(K_n) = 2$. □

Theorem 2.6 *For every two integers n and ℓ with $n \geq 2\ell$, $px_{n,\ell}(K_n) = 2$.*

Proof. It is obvious that $px_{n,\ell}(K_n) \geq 2$. To show that the converse inequality, we will provide an edge-coloring of K_n with 2 colors such that there are at least ℓ internally disjoint spanning proper trees in K_n. It is well known that if n is even, then K_n can be factored into $\frac{n}{2}$ Hamiltonian paths; if n is odd, then K_n can be factored into $\frac{n-1}{2}$ Hamiltonian cycles. In conclusion, K_n contains $\lfloor \frac{n}{2} \rfloor$ pairwise edge-disjoint Hamiltonian paths. Thus, for each Hamiltonian path, we alternately color the edges using color 1 and 2 starting with color 1. If there still remains uncolored edges, then color them with color 1. It is easy to see that there exist $\lfloor \frac{n}{2} \rfloor \geq \ell$ internally disjoint spanning proper trees. Hence, $px_{n,\ell}(K_n) \leq 2$. This completes the proof. □

Remark: Theorem 2.6 is best possible in the sense of the order of K_n. It follows from Theorem 2.1 that there are at most $\lfloor \frac{n}{2} \rfloor$ internally disjoint trees connecting n vertices of K_n. Thus, $\ell \leq \lfloor \frac{n}{2} \rfloor$.

These results naturally lead to the following question: for general integers k, ℓ with $3 \leq k \leq n-1$, whether there exists a positive integer $N_1 = N_1(k, \ell)$ such that $px_{k,\ell}(K_n) = 2$ for every integer $n \geq N_1$? Now, we use two distinct methods to answer this question. Although the result of Theorem 2.9 is stronger than the result of Theorem 2.10, we include both as a demonstration of the variety of possible approaches to this kind of question.

In order to prove Theorem 2.9, we need the following result of Sauer as presented in [3].

Theorem 2.7 *[24] Given $\delta \geq 3$ and $g \geq 3$, for any*

$$m \geq \frac{(\delta-1)^{g-1}-1}{\delta-2},$$

there exists a δ-regular graph G of order $2m$ with girth at least g.

By removing a vertex from the graph provided by Theorem 2.7, we obtain an almost regular graph of odd order, still with girth at least g, but the degree of some vertices is $\delta - 1$. Thus, we replace δ with $\delta + 1$ in Theorem 2.7 to obtain the following easy corollary.

Corollary 2.8 *Given $\delta \geq 3$ and $g \geq 3$, for any*

$$n \geq 2\frac{\delta^{g-1}-1}{\delta-1} - 1,$$

there exists a graph G of order n with $\delta(G) \geq \delta$ and girth at least g.

Theorem 2.9 *Let $k \geq 3$ and $\ell \geq 1$. For all n with*

$$n \geq 2\frac{(\ell(k-1)+k)^4-1}{(\ell+1)(k-1)} - 1,$$

we have $px_{k,\ell}(K_n) = 2$.

Proof. First note that a proper tree using only two colors must be a path. This means that the goal of this result is to produce an edge-coloring with two colors of a complete graph in which any set of k vertices is contained in ℓ internally disjoint proper paths.

By Corollary 2.8 with $g = 5$ and $\delta = \ell(k-1)+k$, we have that for any

$$n \geq 2\frac{(\ell(k-1)+k)^4 - 1}{(\ell+1)(k-1)} - 1,$$

there exists a graph H with n vertices, girth at least 5, and minimum degree at least $\ell(k-1)+k$. Color the edges of H red and color the complement of H blue to complete a coloring of $G = K_n$.

Let S be any set of k vertices in this graph, say with $S = \{v_1, v_2, \ldots, v_k\}$. We say that a path P is S-alternating if each odd vertex of P is in S while each even vertex of P is in $G \setminus S$. Suppose that we have constructed $t \geq 0$ proper paths that are S-alternating, each contain all of S, and are vertex-disjoint aside from the vertices of S. If $t = \ell$, this is the desired system of paths so suppose $t < \ell$ and choose the constructed set of paths so that t is as large as possible.

Further suppose we have constructed an additional S-alternating proper path P^i from v_1 to v_i using red edges of the form $v_j w_j$ for each $j < i$ where $w_j \in G \setminus S$ and blue edges of the form $w_j v_{j+1}$. If $i = k$, this contradicts the choice of t so suppose $i < k$ and further choose this constructed path so that i is as large as possible. Note that at most $\ell(k-1)-1$ vertices of $G \setminus S$ have been used in these existing paths.

Incident to v_i, there are at least $\delta-(\ell(k-1)-1)-(k-1) \geq 2$ red edges to vertices of $G \setminus S$ that have not already been used in paths. Let x and y be the opposite ends of these edges. At most one of x and y, say x, may have a red edge to v_{i+1} since the red graph was constructed to have girth 5. This means that the proper path P^i can be extended to P^{i+1} by including the red edge $v_i y$ and the blue edge $y v_{i+1}$. This contradiction completes the proof. □

Next, we answer the above question by using probabilistic method.

Theorem 2.10 *Let $k \geq 3$ and $\ell \geq 1$. For all n with*

$$n \geq 2k(k+\ell)\ln\left(\frac{1}{1-(1/2)^{2k-3}}\right),$$

we have $px_{k,\ell}(K_n) = 2$.

Proof. Obviously, $px_{k,\ell}(K_n) \geq 2$. For the converse, we color the edges of K_n with two colors uniformly at random. For a k-subset S of $V(K_n)$, let A_S be the event that there exist at least ℓ internally disjoint proper S-trees. Note that a proper tree using only two colors must be a path. It is sufficient to show that $Pr[\bigcap_S A_S] > 0$.

Let S be any k-subset of $V(K_n)$, without loss of generality, we assume $S = \{v_1, v_2, \ldots, v_k\}$. For any $(k-1)$-subset T of $V(K_n)\setminus S$, suppose $T = \{u_1, u_2, \ldots, u_{k-1}\}$,

define $P_T = v_1u_1v_2u_2\cdots v_{k-1}u_{k-1}v_k$ as a path of length $2k-2$ from v_1 to v_k, which implies P_T is an S-tree. Note that for $T, T' \subseteq V(K_n) \setminus S$ and $T \cap T' = \emptyset$, P_T and $P_{T'}$ are two internally disjoint S-trees. Let $\mathcal{P} = \{P_T \mid T \subseteq V(K_n) \setminus S\}$. Take $\mathcal{P}'$ to be a subset of $\mathcal{P}$ which consists of $\lfloor \frac{n-k}{k-1} \rfloor$ internally disjoint S-trees in $\mathcal{P}$. Set $p = Pr[\ P_T \in \mathcal{P}' \text{ is a proper } S\text{-tree }] = \frac{2}{2^{2k-2}} = \frac{1}{2^{2k-3}}$. Let A'_S be the event that there exist at most $\ell-1$ internally disjoint proper S-trees in $\mathcal{P}'$. Assume that $\lfloor \frac{n-k}{k-1} \rfloor > \ell-1$ (that is, $n \geq k + (k-1)\ell$), we have

$$\begin{aligned} Pr[\ \overline{A_S}\ ] &\leq Pr[\ A'_S\ ] \\ &\leq \binom{\lfloor \frac{n-k}{k-1} \rfloor}{\lfloor \frac{n-k}{k-1} \rfloor - (\ell-1)} (1-p)^{\lfloor \frac{n-k}{k-1} \rfloor - (\ell-1)} \\ &= \binom{\lfloor \frac{n-k}{k-1} \rfloor}{\ell-1} (1-p)^{\lfloor \frac{n-k}{k-1} \rfloor - (\ell-1)}. \end{aligned}$$

Then over all possible choices of S with $|S| = k$, we get

$$\begin{aligned} Pr[\ \bigcap_S A_S\ ] &= 1 - Pr[\ \bigcup \overline{A_S}\ ] \\ &\geq 1 - \sum_S Pr[\overline{A_S}] \\ &> 1 - \binom{n}{k} \binom{\lfloor \frac{n-k}{k-1} \rfloor}{\ell-1} (1-p)^{\lfloor \frac{n-k}{k-1} \rfloor - (\ell-1)} \\ &> 1 - n^k \left\lfloor \frac{n-k}{k-1} \right\rfloor^{\ell-1} (1-p)^{\lfloor \frac{n-k}{k-1} \rfloor - \ell + 1} \\ &> 0 \end{aligned}$$

for

$$n \geq 2k(k+\ell) \ln \left(\frac{1}{1-(1/2)^{2k-3}} \right).$$

□

3 Complete bipartite graphs

In this section, we will turn to study the (k, ℓ)-proper index of complete bipartite graphs.

Theorem 3.1 *Let k and ℓ be two positive integers with $k \geq 3$. Then there exists a positive integer N_2 such that $px_{k,\ell}(K_{n,n}) = 2$ for every integer $n \geq N_2$.*

Proof. Clearly, $px_{k,\ell}(K_{n,n}) \geq 2$. We only need to show that $px_{k,\ell}(K_{n,n}) \leq 2$. Firstly, color the edges of $K_{n,n}$ with two colors uniformly at random. For a k-subset S of $V(K_{n,n})$, let B_S denote the event that there exist at least ℓ internally disjoint proper S-trees. Note that a proper tree using only two colors must be a path. It is sufficient to show that $Pr[\bigcap_S B_S] > 0$.

Assume that $K_{n,n} = G[X, Y]$, where $X = \{x_1, x_2, \ldots, x_n\}$ and $Y = \{y_1, y_2, \ldots, y_n\}$. We distinguish the following two cases.

Case 1. Fix the vertices in S in the same class of $K_{n,n}$.

Without loss of generality, we suppose that $S = \{x_1, x_2, \ldots, x_k\} \subseteq X$. Let T be any $(k-1)$-subset of Y, assume that $T = \{y_1, y_2, \ldots, y_{k-1}\}$, define $P_T = x_1y_1 \ldots x_{k-1}y_{k-1}x_k$ as a path of length $2k-2$ from x_1 to x_k, which follows that P_T is an S- tree. Note that for $T, T' \subseteq Y$ and $T \cap T' = \emptyset$, P_T and $P_{T'}$ are two internally disjoint S-trees. Let $\mathcal{P}_1 = \{P_T \mid T \subseteq V(K_n) \setminus S\}$. Take $\mathcal{P}_1'$ to be a subset of $\mathcal{P}_1$ which consists of $\lfloor \frac{n}{k-1} \rfloor$ internally disjoint S-trees in $\mathcal{P}_1$. Set $p_1 = Pr[\ P_T \in \mathcal{P}_1' \text{ is a proper } S\text{-tree }] = \frac{2}{2^{2k-2}} = \frac{1}{2^{2k-3}}$. Let B_S' be the event that there exist at most $\ell - 1$ internally disjoint proper S-trees in $\mathcal{P}_1'$. Assume that $\lfloor \frac{n}{k-1} \rfloor > \ell - 1$ (that is, $n \geq (k-1)\ell$), we have

$$\begin{aligned} Pr[\ \overline{B_S}\ ] &\leq Pr[\ B_S'\ ] \\ &\leq \binom{\lfloor \frac{n}{k-1} \rfloor}{\lfloor \frac{n}{k-1} \rfloor - (\ell-1)} (1-p_1)^{\lfloor \frac{n}{k-1} \rfloor - (\ell-1)} \\ &= \binom{\lfloor \frac{n}{k-1} \rfloor}{\ell-1} (1-p_1)^{\lfloor \frac{n}{k-1} \rfloor - (\ell-1)} \\ &< (\frac{n}{2})^{\ell-1} (1-p_1)^{\lfloor \frac{n}{k-1} \rfloor - (\ell-1)}. \end{aligned}$$

Case 2. Fix the vertices in S in different classes of $K_{n,n}$.

Suppose that $S \cap X = \{s_1, s_2, \ldots, s_r\}$ and $S \cap Y = \{s_{r+1}, s_{r+2}, \ldots, s_k\}$, where r is an positive integer with $1 \leq r \leq k-1$. For any $(k-r)$-subset T_X of $X \setminus S$ with $T_X = \{x_1, x_2, \ldots, x_{k-r}\}$ and any r-subset T_Y of $Y \setminus S$ with $T_Y = \{y_1, y_2, \ldots, y_r\}$, let $T = T_X \cup T_Y$, and define $P_T = s_1y_1 \ldots s_ry_rx_1s_{r+1}x_2 \ldots s_{k-1}x_{k-r}s_k$ as a path of length $2k-1$ from s_1 to s_k. Obviously, P_T is an S-tree. Let $\mathcal{P}_2 = \{P_T \mid T = T_X \cup T_Y, T_X \subseteq X \setminus S, T_Y \subseteq Y \setminus S\}$. Then $\mathcal{P}_2$ has $t = \min\{\lfloor \frac{n-r}{k-r} \rfloor, \lfloor \frac{n-(k-r)}{r} \rfloor\}$ internally disjoint S-trees. Take $\mathcal{P}_2'$ to be a subset of $\mathcal{P}_2$ which consists of t internally disjoint S-trees in $\mathcal{P}_2$. Set $p_2 = Pr[\ P_T \in \mathcal{P}_2' \text{ is a proper } S\text{-tree }] = \frac{2}{2^{2k-1}} = \frac{1}{2^{2k-2}}$. Let B_S'' be the event that there exist at most $\ell - 1$ internally disjoint proper S-trees in $\mathcal{P}_2'$. Note that $\lfloor \frac{n-1}{k-1} \rfloor \leq t < \frac{n}{2}$.

Here, we assume that $\lfloor \frac{n-1}{k-1} \rfloor > \ell - 1$ (that is, $n \geq (k-1)\ell + 1$), we obtain

$$\begin{aligned} Pr[\ \overline{B_S}\ ] &\leq Pr[\ B''_S\ ] \\ &\leq \binom{t}{t-(\ell-1)}(1-p_2)^{t-(\ell-1)} \\ &= \binom{t}{\ell-1}(1-p_2)^{t-(\ell-1)} \\ &< (\frac{n}{2})^{\ell-1}(1-p_2)^{\lfloor \frac{n-1}{k-1} \rfloor-(\ell-1)}. \end{aligned}$$

Since $p_2 < p_1$ and $\lfloor \frac{n-1}{k-1} \rfloor \leq \lfloor \frac{n}{k-1} \rfloor$, we get $Pr[\ \overline{B_S}\ ] < (\frac{n}{2})^{\ell-1}(1-p_2)^{\lfloor \frac{n-1}{k-1} \rfloor-(\ell-1)}$ for every k-subset S of $V(K_{n,n})$. It yields that

$$\begin{aligned} Pr[\ \bigcap_S B_S\ ] &= 1 - Pr[\ \bigcup \overline{B_S}\ ] \\ &\geq 1 - \sum_S Pr[\overline{B_S}] \\ &> 1 - \binom{2n}{k}(\frac{n}{2})^{\ell-1}(1-p_2)^{\lfloor \frac{n-1}{k-1} \rfloor-(\ell-1)} \\ &> 1 - 2^{k-\ell+1}n^{k+\ell-1}(1-p_2)^{\lfloor \frac{n-1}{k-1} \rfloor-(\ell-1)}. \end{aligned}$$

We shall be guided by the above inequality in search for the value of N_2, we find that the inequality $2^{k-\ell+1}n^{k+\ell-1}(1-p_2)^{\lfloor \frac{n-1}{k-1} \rfloor-(\ell-1)} \leq 1$ will lead to $Pr[\ \bigcap_S B_S\ ] > 0$. With similar arguments in Theorem 2.10, we can obtain that there exists a positive integer N_2 for every integer $n \geq N_2$. □

With arguments similar to Theorem 3.1, we can extend the above result to more general complete bipartite graph $K_{m,n}$, where $m = O(n^r)$, $r \in \mathbb{R}$ and $r \geq 1$.

Theorem 3.2 *Let m and n be two positive integers with $m = O(n^r)$, $r \in \mathbb{R}$ and $r \geq 1$. For every pair of integer of k, ℓ with $k \geq 3$, there exists a positive integer $N_3 = N_3(k, \ell)$ such that $px_{k,\ell}(K_{m,n}) = 2$ for every integer $n \geq N_3$.*

4 Random graphs

At the beginning of this section, we introduce some basic definitions about random graphs. The most frequently occurring probability model of random graphs is the Erdös-Rényi random graphs model [16]. The model $G_{n,p}$ consists of all graphs on n vertices in which the edges are chosen independently and randomly with probability p. We say that an event $\mathcal{A}$ happens almost surely if $Pr[\mathcal{A}] \to 1$ as $n \to \infty$.

We now focus on the (k,ℓ)-proper index of the random graph $G_{n,p}$. In what follows, we first show two lemmas which are useful in the main result of this section.

Lemma 4.1 *For any $c \geq 5$, if $p \geq c\sqrt[k]{\frac{\log_a n}{n}}$, then almost surely any k vertices in $G_{n,p}$ have at least $2k^2 \log_a n$ common neighbours, where $a = 1 + \frac{1}{2^{2k-3}-1}$.*

Proof. For a k-subset S of $V(G_{n,p})$, let C_S be the event that all the vertices in S have at least $2k^2 \log_a n$ common neighbours. It is sufficient to prove that for $p = c\sqrt[k]{\frac{\log_a n}{n}}$, $Pr[\bigcap_S C_S] \to 1, n \to \infty$. Let C_1 be the number of common neighbours of all the vertices in S. Then $C_1 \sim B\left(n-k, \left(c\sqrt[k]{\frac{\log_a n}{n}}\right)^k\right)$, and $E(C_1) = \frac{n-k}{n}c^k \log_a n$. In order to apply the Chernoff bound [18] as follows, setting $n > \frac{kc^k}{c^k-2k^2}$.

By the Chernoff Bound, we obtain

$$\begin{aligned}
Pr[\overline{C_S}] &= Pr[C_1 < 2k^2 \log_a n] \\
&= Pr[C_1 < E(C_1)\left(1 - \frac{E(C_1) - 2k^2 \log_a n}{E(C_1)}\right)] \\
&= Pr[C_1 < \frac{n-k}{n}c^k \log_a n \left(1 - \frac{(n-k)c^k - 2k^2 n}{(n-k)c^k}\right)] \\
&\leq e^{-\frac{n-k}{2n}c^k \log_a n\left(\frac{(n-k)c^k-2k^2n}{(n-k)c^k}\right)^2} \\
&< n^{-\frac{c^k(n-k)}{2n}\left(\frac{(n-k)c^k-2k^2n}{(n-k)c^k}\right)^2}.
\end{aligned}$$

Since $1 < a < e$, we have $\log_a n > \ln n$, this leads to the last inequality.

As an immediate consequence, we get

$$\begin{aligned}
Pr[\bigcap_S C_S] &= 1 - Pr[\bigcup_S \overline{C_S}] \\
&\geq 1 - \sum_S Pr[\overline{C_S}] \\
&> 1 - \binom{n}{k} n^{-\frac{c^k(n-k)}{2n}(\frac{(n-k)c^k-2k^2n}{(n-k)c^k})^2} \\
&> 1 - n^{k-\frac{c^k(n-k)}{2n}(\frac{(n-k)c^k-2k^2n}{(n-k)c^k})^2}.
\end{aligned}$$

Note that for any $c \geq 5$, $k - \frac{c^k(n-k)}{2n}(\frac{(n-k)c^k-2k^2n}{(n-k)c^k})^2 < 0$ holds for sufficiently large n. Thus, $\lim_{n\to\infty} Pr[\bigcap_S C_S] = \lim_{n\to\infty} 1 - n^{k-\frac{c^k(n-k)}{2n}(\frac{(n-k)c^k-2k^2n}{(n-k)c^k})^2} = 1$. □

Lemma 4.2 *Let $a = 1 + \frac{1}{2^{2k-3}-1}$. If any k vertices in $G_{n,p}$ have at least $2k^2 \log_a n$ common neighbours, then $px_{k,\ell}(G_{n,p}) \leq 2$ holds almost surely.*

Proof. Firstly, we color the edges of $G_{n,p}$ with two colors uniformly at random. For a k-subset S of $V(G_{n,p})$, let D_S be the event that there exist at least ℓ internally disjoint proper S-trees. Note that a proper tree using only two colors must be a path. If $Pr[\bigcap_S D_S] > 0$, then a suitable coloring of $G_{n,p}$ with 2 colors exists, which follows that $px_{k,\ell}(G_{n,p}) \leq 2$.

We assume that $S = \{v_1, v_2, \ldots, v_k\} \subseteq V(K_n)$, let $N(S)$ be the set of common neighbours of all vertices in S. Let T be any $(k-1)$-subset of $N(S)$, without loss of generality, suppose $T = \{u_1, u_2, \ldots, u_{k-1}\}$, define $P_T = v_1u_1v_2u_2 \cdots v_{k-1}u_{k-1}v_k$ as a path of length $2k-2$ from v_1 to v_k. Obviously, P_T is an S-tree. Let $\mathcal{P}^\star = \{P_T \mid T \subseteq N(S)\}$, then $\mathcal{P}^\star$ has at least $\lfloor \frac{2k^2 \log_a n}{k-1} \rfloor \geq 2k \log_a n$ internally disjoint S-trees (we may and will assume that $2k \log_a n$ is an integer). Take $\mathcal{P}_1^\star$ to be a set of $2k \log_a n$ internally disjoint S-trees of $\mathcal{P}^\star$. It is easy to check that $q = \Pr[\, P_T \in \mathcal{P}_1^\star$ is a proper S-tree $] = \frac{2}{2^{2k-2}} = \frac{1}{2^{2k-3}}$. So $1 - q = a^{-1}$. Let D_1 be the number of proper S-trees in $\mathcal{P}_1^\star$. Then we get

$$\begin{aligned}
Pr[\, \overline{D_S} \,] &\leq Pr[\, D_1 \leq \ell - 1 \,] \\
&\leq \binom{2k \log_a n}{2k \log_a n - (\ell - 1)} (1-q)^{2k \log_a n - (\ell - 1)} \\
&= \binom{2k \log_a n}{\ell - 1} (1-q)^{2k \log_a n - (\ell - 1)} \\
&< (2k \log_a n)^{\ell - 1} a^{-(2k \log_a n - (\ell - 1))} \\
&= \frac{(2ak \log_a n)^{\ell - 1}}{n^{2k}}.
\end{aligned}$$

Consequently

$$\begin{aligned}
Pr[\bigcap_S D_S] &= 1 - Pr[\bigcup \overline{D_S}] \\
&\geq 1 - \sum_S Pr[\, \overline{D_S} \,] \\
&\geq 1 - \binom{n}{k} \frac{(2ak \log_a n)^{\ell - 1}}{n^{2k}} \\
&> 1 - \frac{(2ak \log_a n)^{\ell - 1}}{n^k}.
\end{aligned}$$

It is easy to verify that $\lim_{n \to \infty} 1 - \frac{(2ak \log_a n)^{\ell - 1}}{n^k} = 1$, which implies that $\lim_{n \to \infty} Pr[\bigcap_S D_S] = 1$, this is to say that $px_{k,\ell}(G_{n,p}) \leq 2$ holds almost surely. This completes the proof. □

Combining with Lemmas 4.1 and 4.2, we get the following conclusion.

Theorem 4.3 *Let* $a = 1+\frac{1}{2^{2k-3}-1}$ *and* $c \geq 5$. *For every* $p \geq c\sqrt[k]{\frac{\log_a n}{n}}$, $px_{k,\ell}(G_{n,p}) \leq 2$ *holds almost surely.*

References

[1] N. Alon, J.H. Spencer, *The Probabilistic Method*, John Wiley &Sons, 2004.

[2] E. Andrews, E. Laforge, C. Lumduanhom, P. Zhang, On proper-path colorings in graphs, *J. Combin. Math. Combin. Comput.*, to appear.

[3] B. Bollobás, *Extremal graph theory*, volume 11 of London Mathematical Society Monographs. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1978.

[4] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.

[5] V. Borozan, S. Fujita, A. Gerek, C. Magnant, Y. Manoussakis, L. Montero, Z. Tuza, Proper connection of graphs, *Discrete Math.* **312** (2012) 2550–2560.

[6] Q. Cai, X. Li, J. Song, Solutions to conjectures on (k, ℓ)-rainbow index of complete graphs, *Networks* **62** (2013) 220–224.

[7] Q. Cai, X. Li, J. Song, The (k, ℓ)-rainbow index of complete bipartite graphs, *Bull. Malays. Math. Sci. Soc.*, in press: 10.1007/s40840-016-0348-9.

[8] Q. Cai, X. Li, J. Song, The (k, ℓ)-rainbow index of random graphs, *Bull. Malays. Math. Sci. Soc.* **39(2)** (2016) 765–771.

[9] Q. Cai, X. Li, Y. Zhao, The 3-rainbow index and connected dominating sets, *J. Comb. Optim.* **31(2)** (2016) 1142–1159.

[10] Q. Cai, X. Li, Y. Zhao, Note on the upper bound of the rianbow index of a graph, *Discrete Appl. Math.* **209** (2016) 68–74.

[11] H. Chang, X. Li, Z. Qin, Some upper bounds for the 3-proper index of graphs, arXiv:1603.07840.

[12] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang, The rainbow connectivity of a graph, *Networks* **54** (2009) 75–81.

[13] G. Chartrand, F. Okamoto, P. Zhang, Rainbow trees in graphs and generalized connectivity, *Networks* **55** (2010) 360–367.

[14] L. Chen, X. Li, J. Liu, The k-proper index of graphs, arXiv:1601.06236.

[15] L. Chen, X. Li, K. Yang, Y. Zhao, The 3-rainbow index of a graph, *Discuss. Math. Graph Theory* **35** (2015) 81–94.

[16] P. Erdös, A. Rényi, On the evolution of random graphs, *Publ. Math. Inst. Hungar. Acad. Sci.* **5** (1960) 17–61.

[17] R. Gu, X. Li, Z. Qin, Proper connection number of random graphs, *Theoret. Comput. Sci.* **609(2)** (2016) 336–343.

[18] S. Janson, T. Luczak, A. Ruciński, "Random Graphs " *Whiley-Interscience Series in Discrete Mathesmatics and Optimization,* New York, 2000, xii+333pp.

[19] X. Li, C. Magnant, Properly colored notions of connectivity - a dynamic survey, *Theory & Appl. Graphs* **0**(1) (2015), Art. 2.

[20] X. Li, Y. Mao, *Generalized Connectivity of Graphs*, Springer Briefs in math., Springer, New York, 2016.

[21] X. Li, Y. Shi, Y. Sun, Rainbow connections of graphs: A survey, *Graphs &Combin.* **29** (2013) 1–38.

[22] X. Li, Y. Sun, *Rainbow Connections of Graphs*, Springer Briefs in math., Springer, New York, 2012.

[23] X. Li, M. Wei, J. Yue, Proper connection number and connected dominating sets, *Theoret. Comput. Sci.* **607** (2015) 480–487.

[24] N. Sauer, Extremaleigenschaften regulärer graphen gegebener taillenweite I & II, *Sitzungsberichte Österreich. Akad. Wiss. Math. Natur. Kl.*, S-B II, **176** (1967) 9–25, 27–43.